

# Biomass production of hybrid aspen growing on former farm land in Sweden

Tord Johansson

Received: 2011-11-07; Accepted: 2012-01-13

© Northeast Forestry University and Springer-Verlag Berlin Heidelberg 2012

**Abstract:** We construct dry weight equations for hybrid aspen growing on former farmland in Sweden. Dry weight equations for fractions of hybrid aspen trees were also made. We estimated biomass production in 24 stands. The stands were located in Sweden at latitudes ranging from 55 to 60° N. The mean age was 18 years (range 15–23), the mean stand density 1090 stems·ha<sup>-1</sup> (range 378–2374), and the mean diameter at breast height (over bark) 178 mm (range 85–244 mm). Soil types in the hybrid aspen stands were mainly clay (21 stands), tills (2 stands) and other (1 stand). The mean total standing dry weight above stump level ( $\approx$  200 mm) for the hybrid aspen stands was 135 $\pm$ 53 t·ha<sup>-1</sup> with a range of 42–219 t·ha<sup>-1</sup>. In addition to estimating conventional dry weights of trees and tree components, basic density, specific leaf area (SLA), projected leaf area (PLA) and leaf area index (LAI) were estimated and were in agreement with published figures.

**Keywords:** basic density; biomass production; farmland; hybrid aspen; MAI; plantation forestry

## Introduction

Hybrid aspen has been planted on forest and farm land and studied in Sweden, Finland, Norway, Denmark, Baltic States and in the Great Lake Region in US (Benson and Einspahr 1967; Li and Wu 1998; Dickman 2001; Yu et al. 2001; Rytter and Stener 2003; Karačić et al. 2003; Tullus et al. 2012). Hybridization between the European aspen (*Populus tremula* L.) and the American aspen (*Populus tremuloides* Michx.) was first described at the beginning of the 1920s in Germany (Wettstein 1933). In Sweden the first crossings between *Populus tremula* L. and *Populus*

*tremuloides* Michx. were made in 1939 (Rytter and Stener 2005). Plantations of hybrid aspen even on farmland must be fenced against wild habitat (moose (*Alces alces* L.), deer (*Cervus elaphus* L.), roedeer (*Capreolus capreolus capreolus* L.), hare (*Lepus capensis* L., *Lepus timidus* L.), rabbit (*Oryctolagus cuniculus* Pallas) and vole (*Microtus agrestis* L., *Arvicola amphibious* L.)). Plantations on forest land in Sweden are sparse and not recommended depending on the high wild habitat population. There are some Nordic results from hybrid aspen trials established in the period 1940 to 1960 (Johansson 1953, 1976; Langhammer 1973; Jacobsen 1976; Eriksson 1984). Mean annual increment (MAI) for the measured trials varied between 12 and 17 m<sup>3</sup>·ha<sup>-1</sup>·a<sup>-1</sup> ( $\approx$ 4.2–6.0 t·ha<sup>-1</sup>·a<sup>-1</sup>). An advantage with hybrid aspen as well as the parent species is the rapid establishment and fast growth of the second generation. Hybrid aspen produces 50,000 to 100,000 suckers after cutting (Rytter 2006). In a German study the number of suckers per hectare established after harvesting the first plantation (10 years) was 165,000 and after the second rotation (10 years) the number was 215,000 (Liese-bach et al. 1999).

During the last 20 years, there has been a general increase in interest in the management of broadleaved trees in the Nordic countries (and elsewhere), despite the lack of suitable climate-adapted clones and uncertainties regarding appropriate management, pest control, economic factors, markets and future land-use policy at both EU and national levels etc. The earliest cultivation of hybrid aspen in Sweden started in the end of the 1930s (Johansson 1953, 1976; Langhammer 1973). In the late 1940s the Swedish Match Company showed strong interest in hybrid aspen as a fast growing and high yielding species. A breeding program was started mainly focused on fast growing individuals with a high yield (Rytter and Stener 2005). Later on in the 1980 the objective for hybrid aspen cultivation changed to focusing on the specie as a crop on former farm land. Today there is an increasing interest in biofuel as an energy supply. In Sweden the main biofuel sources are tops and twigs collected after clear cuttings. Among other crops useable as biofuel hybrid aspen is important specie to test (Karlsson 1991; Christersson 1996; Ilstedt and Gullberg 1993). In Sweden hybrid aspen is

---

The online version is available at <http://www.springerlink.com>

Tord Johansson

Department of Energy and Technology, Swedish University of Agricultural Sciences, 750 07 Uppsala, SWEDEN.

E-mail: tord.johansson@slu.se

---

Responsible editor: Chai Ruihai

mainly used as pulpwood, wood for matches and in a small scale as plywood. These scenarios might be an interesting alternative both on farmland and on fertile forest land. Then with the use of hybrid aspen wood for different purposes in Norway and Finland there are existing markets for hybrid aspen pulpwood, saw logs and biofuels. Few studies dealing with biomass yield from hybrid aspen stands growing on farmland are published.

The present study is a part of a series on the biomass of broad-leaf species, either planted or self-generated, and growing on forest or farm land (Johansson 1996, 1999a, 1999b, 1999c, 1999d, 2000, 2002, 2006, 2008, 2011; Johansson and Karačić 2011). Since reports on leaf characteristics for hybrid aspen, such as projected leaf area (PLA), leaf area index (LAI) and specific leaf area (SLA), are sparse but important structural parameters of forest ecosystems, leaf and canopy measurements were made. Leaf characteristics, mainly LAI, have an important influence on the exchange of energy, gas and water in trees. LAI is a key component of biochemical cycles in ecosystems (Bréda 2003).

The aim of the present study was to construct dry weight equations for hybrid aspens and their above-ground components. Constructed biomass estimation functions were used for calculating biomass production for plantations on former farmland. The effect of age and density on biomass production was analyzed by the constructed biomass function. Characteristics such as LAI, SLA and PLA were estimated and calculated for both individual trees and entire stands.

## Material and methods


### Study site

In this study, we evaluated 24 stands of hybrid aspen growing on former farmland. The study area is located between latitudes 56° and 60° N in Sweden (Fig. 1 and Table 1). No information about clones was available but the material was a hybrid between European aspen origin lat. 56–57° N. and American aspen origin lat. 43–54° N. However later on in 2000 it has been introduced improved clones c.f. Stener and Karlsson (2004). The age of the stands ranged from 15 to 23 years and the stem number ranged from 378 to 2374 (Table 1). Initially 1100–2500 stems per hectare was planted. Thinning operations have been made only in stands no. 3, 13 and 23. The thinning removals were 50% of initial stem number (1500, 2500 and 1100 respectively). The low stem numbers in some of the studied stands indicate loss of stems caused by among others wild habitat. As the stands were established on farmland no other tree species than hybrid aspen were growing on the site. Early growth and damage to the plantation were assessed on the basis of information provided by the forest owner.

### Sampling and measurements

The mean number of stems per hectare was calculated based on the number of stems counted in either whole stands or plots, as follows. The area of the studied plantations varied between 0.09

and 2.40 ha. In stands > 1 ha stands, a 0.5 ha plot in the central part of each stand was chosen: The outer row in the stand was not included avoiding edge effects caused by factors such as wind, open areas, ditches and shading by adjacent stands). The diameter at breast height (DBH) of each counted tree was measured by cross callipering (Table 1), and the arithmetic mean diameter was calculated for each stand.



**Fig. 1** Localities of sample trees of hybrid aspen growing on abandoned farmland in Sweden.

For calculation of mean height in the hybrid aspen stands a statistical regression model for each stand was made. The height and diameter for 9 trees in two subsample plots each was measured. The two subsample plots (3 trees by 3 rows close to each other) were systematically located e.g. the first measured tree was the fifth tree in the third row. The other subsample plots started in the opposite part of the stand area diagonal to the first plot. Then a regression model for each stand describing the relation of height to diameter was made. Tree mean height on the plots was calculated by the regression models.

In each stand, one hybrid aspen with a diameter close to the estimated arithmetic mean diameter was selected. This tree was then assessed in terms of damage, stem straightness, and leader type (single vs. double or broken). The tree also had to be free from root rot. If any tree was damaged or rotten, it was not used, and another from the stand was selected instead. The trees were felled and cut as near to the ground as possible and their height (cm) were measured (Table 2) and DBH (mm) was recorded. The age of these trees was determined in the laboratory by counting annual rings in discs taken from the base of the stem (total age) and breast height (breast height age). The stem was weighed fresh in the field. All twigs on the tree were then cut and weighed with leaves. Since the sampling procedure to quantify the leaf mass is very laborious, a "mean leaf technique" method was used

for dry mass estimation. All leaves from three hybrid aspens from each of three localities, totally 9 trees, were sampled and weighed fresh. The samples were taken from trees with stem diameters of 100, 200 and 300 mm DBH. Then a simple allomet-

ric equation was derived, relating DBH to percentage fresh leaf weight by fresh weight of twigs + leaves, and used for estimating the leaf weight of all sampled trees.

**Table 1. Main characteristics of sampled hybrid aspen stands growing at 24 locations in Sweden**

| Plot No | Age, years | Height, m<br>mean±SD | DBH, mm<br>mean±SD | No. of stems ha <sup>-1</sup> | Basal area m <sup>2</sup> ·ha <sup>-1</sup> | Soil type        | Stand/plot size, ha |
|---------|------------|----------------------|--------------------|-------------------------------|---|------------------|---------------------|
| 1       | 19         | 11.8±2.9             | 126±62             | 850                           | 10,6  | Sandy silty till | 0.09/0.08           |
| 2       | 18         | 22.5±2.1             | 191±39             | 1467                          | 42,0  | Light clay       | 0.11/0.09           |
| 3       | 19         | 25.8±2.0             | 224±32             | 710                           | 28,0  | Light clay       | 0.42/0.40           |
| 4       | 19         | 20.8±1.5             | 237±42             | 432                           | 19,0  | Light clay       | 0.08/0.07           |
| 5       | 19         | 18.3±3.5             | 167±40             | 814                           | 17,8  | Light clay       | 0.08/0.07           |
| 6       | 19         | 14.9±3.7             | 140±44             | 650                           | 10,0  | Fine sand        | 0.09/0.08           |
| 7       | 17         | 19.7±2.8             | 158±44             | 1540                          | 30,2  | Medium clay      | 0.09/0.07           |
| 8       | 17         | 17.1±2.5             | 125±26             | 2232                          | 27,4  | Light clay       | 0.19/0.18           |
| 9       | 17         | 16.5±1.6             | 157±34             | 844                           | 16,3  | Medium clay      | 0.10/0.09           |
| 10      | 23         | 20.2±1.8             | 197±23             | 1050                          | 32,0  | Light clay       | 0.10/0.08           |
| 11      | 19         | 17.5±3.4             | 173±44             | 1370                          | 32,2  | Light clay       | 0.11/0.10           |
| 12      | 19         | 15.5±3.3             | 147±40             | 1760                          | 29,9  | Light clay       | 0.13/0.10           |
| 13      | 17         | 16.9±2.1             | 161±41             | 1150                          | 23,4  | Light clay       | 2.40/0.50           |
| 14      | 17         | 19.4±1.8             | 235±33             | 486                           | 21,1  | Light clay       | 0.26/0.24           |
| 15      | 20         | 20.6±1.7             | 162±31             | 567                           | 11,7  | Medium clay      | 0.11/0.10           |
| 16      | 18         | 25.0±2.3             | 237±59             | 779                           | 34,3  | Light clay       | 0.12/0.10           |
| 17      | 16         | 10.4±1.4             | 85±26              | 1259                          | 7,1   | Light clay       | 0.09/0.08           |
| 18      | 16         | 14.6±2.0             | 150±46             | 2000                          | 35,3  | Light clay       | 0.09/0.08           |
| 19      | 15         | 16.0±1.8             | 142±25             | 2374                          | 37,6  | Medium clay      | 0.13/0.12           |
| 20      | 18         | 20.8±1.5             | 179±38             | 1600                          | 40,2  | Light clay       | 0.13/0.11           |
| 21      | 19         | 22.6±1.7             | 225±39             | 378                           | 15,0  | Sandy silty till | 0.31/0.30           |
| 22      | 19         | 24.1±2.1             | 239±63             | 468                           | 21,0  | Medium clay      | 0.36/0.35           |
| 23      | 20         | 22.2±1.0             | 244±32             | 593                           | 27,7  | Medium clay      | 0.20/0.18           |
| 24      | 20         | 20.8±2.0             | 181±44             | 785                           | 20,2  | Light clay       | 0.22/0.21           |
| Mean±SD | 18±2       | 18.9 ±3.9            | 178±43             | 1090±585                      | 24.6±10.0                                   |                  |                     |
| Range   | 15–23      | 13.5–25.8            | 85–244             | 378–2374                      | 7.1–42.0                                    |                  |                     |

Samples of stem and twigs were taken for estimating basic wood density. A disc at 4 m height was taken. Four samples of twigs (10 cm long) were taken close to the stem base, two in the upper and two in the lower part of the crown. The basic density of stems and twigs was estimated according to the water-immersion method described by Andersson and Tuimala (1980). Samples of stems and twigs were saturated in water for 24 h and then weighed and their volume (cm<sup>3</sup>) was determined. The dry matter content of the wood proportion (g) of the samples was determined after drying at 105°C in an air-ventilated oven for 3–5 days, depending on their dimensions. Dry weight to fresh volume ratios of the debarked disk and twigs were then calculated as basic density (g·cm<sup>-3</sup>) (Table 2).

A sample of 200 leaves from sample trees in all hybrid aspen stands was taken from the top, middle and lower parts of the crowns then frozen for analyses in the laboratory. The 200 sampled leaves were weighed fresh, and the estimation of PLA (projected leaf area), mm<sup>2</sup>, of the sample was determined with a leaf-area meter (LI-3000, LI-COR, Inc. Lincoln, Neb.). Each of the samples was then dried at 105°C in an oven for 48 h and weighed. The dry mass of the leaves was then calculated. The specific leaf area, SLA (cm<sup>2</sup> g<sup>-1</sup>) was calculated as the total leaf area (cm<sup>2</sup>) of a tree divided by the dry weight (kg) of the leaf

mass.

Within each sample hybrid aspen stand, leaf area index (LAI) was estimated using a LAI-2000 plant canopy analyzer (LI-COR Inc., Lincoln, Neb.). An initial measurement was taken outside the stand then another five readings were taken along each of five transects inside the stand, giving in total 25 estimations from various points within the stand. Measurement locations were both under hybrid aspen trees and in small stand openings. A final measurement was taken outside the stand to calibrate LAI.

Soil samples, down to a depth of 30 cm, were taken from two points in each stand, and the mean texture of the sampled layer was determined. Soils were classified as tills and sediments (Johansson, 1999) in the field, following guidelines provided by Ekström (1926) and then according to particle size in the laboratory. The particle size distribution was determined using a mechanical sieving method (English and German standard), and soil types were classified as follows: sediments as gravel (20–2 mm), coarse sand (2–0.2 mm), fine sand (0.2–0.02 mm), silt (0.02–0.002 mm) or clay (<0.002 mm); tills as gravel, sandy, fine sandy or silty tills; and organogenic soils as moorland peat or moss peat. Although the soil samples contained particles of different size, their type designation was based on the most frequent size, together with one or two prefixes associated with

other less frequent soil types. Clay soils were then classified based on their percentage clay as follows: light clay (13%–29%), medium clay (30%–40 %), heavy clay (41%–60%), and till clay (13%–60%) The water table was estimated to be between 30 cm and 60 cm deep at all sites. Most of the hybrid aspen stands were growing on sediments: 6 on medium clay, 15 on light clay and 3 on tills.

**Table 2. Characteristics of sampled trees of hybrid aspen stands.**

| Locality No | Age, yrs | Height, m | Diameter, DBH, ob | MAI kg yr <sup>-1</sup> | Basic density, g cm <sup>-3</sup> |             |
|-------------|----------|-----------|-------------------|-------------------------|-----------------------------------|-------------|
|             |          |           |                   |                         | Stems                             | Twigs       |
| 1           | 19       | 16.7      | 236               | 13.0                    | 0.320                             | 0.430       |
| 2           | 18       | 20.5      | 223               | 9.4                     | 0.305                             | 0.293       |
| 3           | 19       | 23.0      | 255               | 13.8                    | 0.295                             | 0.298       |
| 4           | 19       | 21.1      | 260               | 10.4                    | 0.371                             | 0.412       |
| 5           | 19       | 15.1      | 221               | 11.0                    | 0.401                             | 0.328       |
| 6           | 19       | 17.3      | 242               | 7.6                     | 0.345                             | 0.312       |
| 7           | 17       | 20.0      | 189               | 7.2                     | 0.343                             | 0.326       |
| 8           | 17       | 20.3      | 145               | 4.0                     | 0.346                             | 0.533       |
| 9           | 17       | 17.1      | 177               | 5.6                     | 0.290                             | 0.309       |
| 10          | 23       | 21.2      | 191               | 6.4                     | 0.314                             | 0.300       |
| 11          | 19       | 21.4      | 206               | 10.8                    | 0.494                             | 0.432       |
| 12          | 19       | 17.5      | 219               | 9.6                     | 0.459                             | 0.361       |
| 13          | 17       | 19.6      | 227               | 10.8                    | 0.338                             | 0.300       |
| 14          | 17       | 18.1      | 270               | 19.4                    | 0.411                             | 0.489       |
| 15          | 20       | 18.0      | 292               | 16.9                    | 0.343                             | 0.290       |
| 16          | 18       | 24.3      | 338               | 18.1                    | 0.361                             | 0.340       |
| 17          | 15       | 14.0      | 149               | 5.1                     | 0.378                             | 0.333       |
| 18          | 15       | 17.0      | 211               | 9.3                     | 0.371                             | 0.319       |
| 19          | 16       | 16.5      | 182               | 8.6                     | 0.343                             | 0.320       |
| 20          | 18       | 21.1      | 242               | 13.4                    | 0.344                             | 0.316       |
| 21          | 19       | 21.9      | 263               | 13.7                    | 0.313                             | 0.333       |
| 22          | 19       | 25.4      | 388               | 32.9                    | 0.356                             | 0.373       |
| 23          | 20       | 23.2      | 310               | 18.6                    | 0.306                             | 0.341       |
| 24          | 20       | 22.0      | 195               | 8.1                     | 0.400                             | 0.610       |
| Mean±SD     | 18±2     | 19.7±2.9  | 235±57            | 11.9±6.0                | 0.356±0.050                       | 0.362±0.083 |
| Range       | 15–23    | 14.0–25.4 | 145–388           | 5.1–32.9                | 0.290–0.494                       | 0.290–0.610 |

## Data analyses

The dry mass production per tree was calculated on the basis of an equation describing the correlation between DBH and dry mass production (kg/tree), derived from data collected from all of the measured trees.

The power function was tested:

$$M = \beta_0 \times D^{\beta_1} \quad (1)$$

where,  $M$  is the dry mass, kg tree<sup>-1</sup>,  $D$  is the diameter at breast height, over bark (ob), mm,  $\beta_0$  and  $\beta_1$  are parameters.

The power model is frequently used to describe such relationships (Kittredge 1944; Payandeh 1981; Satoo and Madgewick 1985; Bolstad and Gower 1990; Johansson 1999). Based on the mean diameter, the actual dry mass production of each of the hybrid aspen stands included in the study was estimated.

Data were analyzed by nonlinear regression using the SAS/STAT system for personal computers (SAS 2006). A meas-

ure of the fit of the nonlinear regressions was based on the coefficient of determination (Zar 1999):

$$R^2 = 1 - (\text{SSE}/\text{SST (corrected)}) \quad (2)$$

where SSE is the sum of squares of the error terms and SST is the total sum of squares.

The regression was also tested by root mean squared error (RMSE).

$$RMSE = 1 - \sum (Y_i - \hat{Y})^2 / (Y - \bar{Y})^2 \quad (3)$$

Throughout the report, means are presented together with standard deviation (SD).

## Results

### Allometric relationships

The tested function (1) fitted the data relating dry mass to DBH for both small and large hybrid aspens and the curves described by the function are statistically acceptable. The value of the determination coefficient ( $R^2$ ) and RMSE indicate good correlation between the fitted curves and the estimated values. Further information about parameter estimates is given in Table 3.


**Table 3. Estimated parameters of Equation (1) for dry weight estimations of hybrid aspen growing on former farmland**

| Components | Parameter | Parameter estimates | Standard errors of parameters | $R^2$ | RMSE  |
|------------|-----------|---------------------|-------------------------------|-------|-------|
| Total      | $B_0$     | 0.00666             | 0.00598                       | 0.97  | 46.35 |
|            | $B_1$     | 1.90220             | 0.15920                       |       |       |
| Stem       | $B_0$     | 0.00273             | 0.00230                       | 0.97  | 31.40 |
|            | $B_1$     | 2.00430             | 0.14880                       |       |       |
| Twigs      | $B_0$     | 0.00881             | 0.00002                       | 0.81  | 28.48 |
|            | $B_1$     | 1.59260             | 0.41510                       |       |       |
| Leaves     | $B_0$     | 0.00227             | 0.00022                       | 0.96  | 1.94  |
|            | $B_1$     | 1.92670             | 0.13530                       |       |       |


### Biomass structure for sample trees

The mean total dry tree weight (range within parenthesis) was 218±120 (68–625) kg (Table 4) and MAI was 11.9±6.0 (5.1–32.9) kg year<sup>-1</sup> (Table 2). Curves relating the dry masses of the various fractions per tree (total dry mass above stump level, stem, twigs and leaves) to diameter are presented in Fig. 2. The mean percentage ± SD of the total dry weight accounted for by the dry weight of stem, twigs and leaves was 72±10 % (51–85), 24±10 % (11–46) and 4±1 (3–7), respectively (Fig. 3 and Table 4). The mean percentage dry matter of fresh weight for stem, twigs and leaves was 54±5 (47–65), 53±5 (46–69) and 37±4 (27–47). Basic density for the 25 sampled hybrid aspen stems varied between 0.295 and 0.494 g·cm<sup>-3</sup> with a mean of

$0.356 \pm 0.050 \text{ g} \cdot \text{cm}^{-3}$  and for twigs  $0.362 \pm 0.083$  ( $0.290$ – $0.610$ ) (Table 2).



**Fig. 2** Dry mass production per tree,  $\text{kg tree}^{-1}$ , in relation to diameter at breast height (DBH), mm of total, stem, twigs and leaves for sample trees of hybrid aspen stands growing on abandoned farmland



**Fig. 3** Percentage of total tree dry weight accounted for by stem, twigs and leaves in relation to diameter at breast height (DBH; mm) for sample trees of hybrid aspens growing on farmland.


**Table 4.** Fresh-weight and dry-mass production ( $\text{kg} \cdot \text{tree}^{-1}$ ) and mean percentage of total tree fresh-weight and dry-weight accounted for by stem, twigs and leaves from sampled hybrid aspen trees

|               | Fresh-weight, $\text{kg} \cdot \text{tree}^{-1}$ |               |              |             | Percentage of total fresh-weight |             |           |
|---------------|--|---------------|--------------|-------------|----------------------------------|-------------|-----------|
|               | Total  | Stem          | Twigs        | Leaves      | Stem                             | Twigs       | Leaves    |
| Mean $\pm$ SD | 411 $\pm$ 225                                    | 286 $\pm$ 163 | 101 $\pm$ 69 | 24 $\pm$ 13 | 70 $\pm$ 10                      | 24 $\pm$ 9  | 6 $\pm$ 1 |
| Range         | 145–1185   | 89–859        | 23–255       | 11–71       | 50–85                            | 7–46        | 4–9       |
|               | Dry-weight, $\text{kg tree}^{-1}$                |               |              |             | Percentage of total dry-weight   |             |           |
|               | Total  | Stem          | Twigs        | Leaves      | Stem                             | Twigs       | Leaves    |
| Mean $\pm$ SD | 218 $\pm$ 120                                    | 157 $\pm$ 90  | 53 $\pm$ 36  | 9 $\pm$ 5   | 72 $\pm$ 10                      | 24 $\pm$ 10 | 4 $\pm$ 1 |
| Range         | 68–625   | 41–467        | 13–132       | 3–26        | 51–85                            | 11–46       | 3–7       |

#### Biomass structure for hybrid aspen stands

Actual dry mass production per hectare for the stands was calculated (Table 5). The mean total dry mass above stump level was

$135 \pm 53 \text{ t} \cdot \text{ha}^{-1}$  (42–219), and the mean MAI was  $7 \pm 3 \text{ t} \cdot \text{ha}^{-1} \cdot \text{a}^{-1}$  (3–13). The correlation ( $R^2=0.16$ ,  $p<0.0001$ ) between total dry mass production above stump level and DBH for hybrid aspen stands is illustrated in Fig. 4. Mean leaf area index (LAI) was  $2.24 \pm 0.58$  (0.68–3.35) (Table 6).



**Fig. 4** Total dry mass production (PROD) above stump level in relation to diameter at breast height for hybrid aspen stands growing on abandoned farmland

**Table 5.** Dry-mass production ( $\text{t} \cdot \text{ha}^{-1}$ ) and mean annual increment (MAI;  $\text{t} \cdot \text{ha}^{-1} \cdot \text{a}^{-1}$ ) of whole hybrid aspen stands.

| Plot no.                       | Total        | Stem        | Twigs       | Leaves    | MAI       |
|--------------------------------|--------------|-------------|-------------|-----------|-----------|
| 1                              | 67.0         | 40.6        | 18.1        | 2.0       | 3.5       |
| 2                              | 219.3        | 161.0       | 60.4        | 8.0       | 12.2      |
| 3                              | 144.0        | 107.3       | 37.7        | 5.3       | 7.6       |
| 4                              | 97.5         | 73.1        | 25.1        | 3.5       | 5.1       |
| 5                              | 96.6         | 68.3        | 27.1        | 3.4       | 5.1       |
| 6                              | 56.6         | 38.3        | 16.3        | 1.9       | 3.0       |
| 7                              | 167.2        | 115.7       | 46.8        | 5.9       | 9.8       |
| 8                              | 151.8        | 104.9       | 46.7        | 5.4       | 8.9       |
| 9                              | 88.7         | 62.6        | 25.4        | 3.1       | 5.2       |
| 10                             | 164.9        | 122.7       | 45.4        | 6.1       | 7.2       |
| 11                             | 174.9        | 123.3       | 48.2        | 6.1       | 9.2       |
| 12                             | 171.6        | 114.3       | 47.8        | 5.8       | 9.0       |
| 13                             | 128.3        | 89.6        | 36.1        | 4.5       | 7.5       |
| 14                             | 106.9        | 80.8        | 27.9        | 3.9       | 6.3       |
| 15                             | 62.4         | 44.7        | 18.0        | 2.2       | 3.1       |
| 16                             | 180.7        | 131.9       | 45.3        | 6.4       | 10.0      |
| 17                             | 42.3         | 27.3        | 14.3        | 1.4       | 2.8       |
| 18                             | 198.4        | 135.4       | 56.0        | 6.8       | 13.2      |
| 19                             | 201.6        | 143.9       | 61.1        | 7.2       | 12.6      |
| 20                             | 214.7        | 154.3       | 59.5        | 7.7       | 11.9      |
| 21                             | 88.5         | 57.6        | 20.2        | 2.8       | 2.8       |
| 22                             | 125.2        | 80.5        | 27.5        | 3.9       | 3.9       |
| 23                             | 164.2        | 106.4       | 36.1        | 5.2       | 5.2       |
| 24                             | 130.3        | 77.4        | 29.7        | 3.9       | 3.9       |
| Mean $\pm$ SD                  | 135 $\pm$ 53 | 94 $\pm$ 37 | 37 $\pm$ 14 | 5 $\pm$ 2 | 7 $\pm$ 3 |
| Range                          | 42–219       | 27–161      | 16–61       | 1–8       | 3–13      |
| Percentage of total dry-weight |              |             |             |           |           |
| Mean $\pm$ SD                  | 70 $\pm$ 2   |             | 27 $\pm$ 2  | 3 $\pm$ 1 |           |
| Range                          | 64–72        |             | 25–33       | 3–4       |           |

**Table 6.** Leaf area index (LAI) in hybrid aspen stands, individual leaf dry-weight (mg) and total leaf weight per tree (g), number of leaves per tree, projected leaf area (PLA; cm<sup>2</sup>) and specific leaf area (SLA; cm<sup>2</sup>·g<sup>-1</sup>) of sample trees.

|         | LAI       | Weight d.w.             |                               | No. of<br>leaves tree <sup>-1</sup> | PLA<br>cm <sup>2</sup> | SLA<br>(cm <sup>2</sup> ·g <sup>-1</sup> d.w.) |
|---------|-----------|-------------------------|-------------------------------|-------------------------------------|------------------------|--|
|         |           | leaf <sup>-1</sup> (mg) | leaves tree <sup>-1</sup> (g) |                                     |                        |  |
| Mean±SD | 2.24±0.58 | 373±245                 | 8324±4806                     | 28306±20463                         | 48.86±13.76            | 165.75±81.11                                   |
| Range   | 0.68–3.35 | 174–1312                | 3500–26400                    | 8197–81589                          | 29.74–71.44            | 33.17–350.93                                   |

## Discussion

Measured values in the present study represent actual dry weight biomass. In the present study, twig and leaf weight as a percentage of total hybrid aspen dry weight decreased with increasing tree diameter (Fig. 3). The mean standing biomass of hybrid aspen of 135±53 t·ha<sup>-1</sup> (42–219), is within the range of other published figures (Table 7). However, in some cases, the results are not comparable, since the data are not presented with diameter as the dependent variable. Another problem is that yield

In the present study, the mean annual increment (MAI) of total dry weight was 7±3 (3–13) t·ha<sup>-1</sup>·a<sup>-1</sup>. These values are similar to those reported in other studies for stands of the same age (Table 7). When comparing the standing dry weight production above stump level with corresponding estimates made in previous studies, it is important to keep in mind that some of the stands in the present study had low numbers of stems depending on damages. Thus, a more relevant comparison is the relationship between DBH and mean single-tree dry weight values based on either whole-tree or tree-component data. On this basis, the dry weight production values obtained for hybrid aspen stands in the present study (Table 4) are similar to, or higher than, those in other reports (Table 7).

Mostly the low number of aspens in the studied stands is depending on the lack of efficient fence against the strong pressure by wild habitat mostly moose and hare. Furthermore the ground vegetation indicated a poor or missing soil treatment before planting. Other factors could be high dryness close to initial plantation time. There are no indications that the biomass production in stands growing in southern Sweden (nos. 21–23) is higher than in other parts of Sweden.

However in Nordic countries there are some other fast-growing broad leaf species, which may be suitable for biomass production. In Table 7 published biomass production of alder (*Alnus glutinosa* (L.) Gaertner; *Alnus incana* (L.) Moench), European aspen (*Populus tremula* L.) and birch (*Betula pendula* Roth; *Betula pubescens* Ehrh.) and hybrid poplar are presented. MAI for those species differ between 1 and 15 t·ha<sup>-1</sup>·a<sup>-1</sup>. Values for 10–30 year-old European aspen, alder and birch stands were: 3–14, 2–10 and 1–8 t·ha<sup>-1</sup>·a<sup>-1</sup> respectively. The stands were naturally regenerated and mostly growing on forest land and not managed. The area of studied stands was small but the number of stems per hectare was high >3000. After harvest natural regeneration by sprouts and suckers might produce as high biomass as the presented figures. MAI for 10–30-year-old hybrid poplar plantation growing on former farmland was 3–15 t·ha<sup>-1</sup>·a<sup>-1</sup>, which is similar to those found for hybrid aspen in the present

figures are presented in volume (m<sup>3</sup>) instead of biomass. As a rough measure the volume figures can be converted to biomass by multiplying by 0.350. The standing biomass per hectare for some stands is big when studying the number of stems (1400–1600) and mean diameter (170–190 mm). However, mean diameters of 160 to 220 mm and corresponding stem numbers 1400–1700 per hectare are noted for European aspen (Johansson 1999). There were no indications of self regeneration neither in the present nor the previous studies. But generally most of these stands require thinning in order to maintain high productivity rate until final harvest.

Hybrid aspen stems accounted for 72±10 (50–85)% of the total aboveground dry biomass in the stands examined in the present study, while twigs accounted for 24±9% and leaves for 4±1%. Similarly, other studies have reported proportions of 73% for stems and 27% for twigs of 7-year old hybrid aspens growing on former farmland in Estonia (Tullus et al. 2009). Rytter and Stener (2003) reported proportions 76% for stems and 24% for twigs and leaves in a 14-year-old hybrid aspen stands.

The basic density indicates the amount of solid wood produced by the tree. This is an important factor when choosing a suitable species or clone for commercial forest management. As shown in the material examined here the mean basic density for stems was 0.356±0.050 (0.295–0.494) g·cm<sup>-3</sup>. The range of the values means that it may be possible to choose clones or species with a high basic density. Similar values have been reported: 0.348 g·cm<sup>-3</sup> for hybrid aspens grown in southern Sweden (Rytter and Stener 2003), 0.380 g·cm<sup>-3</sup> in another study of 26-year-old hybrid aspens (Ilstedt and Gullberg 1993). Elfving 1986a) reported basic density of 0.348 g·cm<sup>-3</sup> for 26-year-old hybrid aspens growing in southern Sweden and in another study 0.355 g·cm<sup>-3</sup> (on average) for 10-year old trees with DBH 60–160 mm growing in southern Sweden (Stener 1998). In two American studies (Einspahr et al. 1968; Einspahr and Wyckoff 1975) basic densities of 0.403 (0.367–481) and 0.410 g·cm<sup>-3</sup> respectively were reported from Lake States.

Generally, the annual production of foliage has been estimated either on the basis of standing biomass or from litter-fall. In the present study, estimates of standing foliage dry weight were based on mean data for the trees with the best conditions for foliage growth in the stand e.g. samples were made in July–August. However, the foliage dry weight can be affected by factors other than tree height and diameter (Madgwick 1971). To understand how the canopy contributes to total photosynthesis, knowledge is needed on how the distribution of foliage varies among dominant, co-dominant and suppressed trees, as the stand develops. In a study by Woods et al. (1991) about statistical error

analysis for biomass and leaf area index estimations for trembling aspen they reported a range of 1.3 to 4.0. According to the report LAI did not show clear relationship with biomass amount, stand density or average stem diameter. These patterns are ex-

pected for an early succession, shade-intolerant species. LAI remained relatively constant once a full canopy was established about 8 years after clear cutting of the studied stands (Woods et al. 1991).

**Table 7. Above-ground biomass production by hybrid aspen trees and stands and other fast-growing broad leaf species in Nordic countries. Data from published literature in chronological order of publication.**

| Stand age, years                                     | Tree weight, kg (mm) | Stand production t·ha <sup>-1</sup> | MAI, t·ha <sup>-1</sup> ·a <sup>-1</sup> | Remarks                  | References                   |
|--|----------------------|-------------------------------------|--|--------------------------|------------------------------|
| 5  | 6.3                  |                                     |  | Finland                  | Yu et al. 2001               |
| 14   | 61.7 (154)           | 98.7                                | 7.1                                      | Sweden                   | Rytter and Stener 2003       |
| 11   |                      | 77.0                                | 7.0                                      | Sweden                   | Karačić et al. 2003          |
| 14   |                      | 42–168.0                            | 3.0–12.0                                 | Sweden                   | Rytter and Stener 2005       |
| 23   |                      | 207.0                               | 7.0                                      | Sweden                   | Johnsson 1953                |
| 16   | (146)                | 120.0                               | 7.5                                      | Norway                   | Langhammer 1973              |
| 30   |                      | 117.0                               | 3.9                                      | Central Sweden           | Eriksson 1984                |
| 28   |                      | 168.0                               | 6.0                                      | Denmark                  | Jacobsen 1976                |
| 10   | 26.9 (87)            | 120.0                               | 12.0                                     | Germany                  | Liesebach et al. 1999        |
| 26   |                      | 119.6–208.0                         | 4.6–8.0                                  | Southern Sweden          | Istedt and Gullberg 1993     |
| 7  | (31)                 | 8.0                                 | 1.1                                      | Estonia                  | Tullus et al. 2009           |
| 8  | 5.0                  | 24.3                                | 3.0                                      | Sweden                   | Telenius 1999                |
| 7–15   |                      | 19.6–76.5                           | 2.8–5.1                                  | Sweden                   | Rytter 2002                  |
| 5  |                      | 26.5                                | 5.4                                      | Rotation no. 1 Germany   | Hofmann-Schielle et al. 1999 |
| 5  |                      | 53.5                                | 10.7                                     | Rotation no. 2 Germany   |                              |
| 24   | (185)                | 100.5                               | 4.2                                      | Sweden                   | Persson 1973                 |
| 42   | (310)                | 198.1                               | 4.5                                      | Sweden                   |                              |
| 26   |                      | 156.8                               | 5.6                                      | Southern Sweden          | Elfving 1986a                |
| 32   |                      | 169.6                               | 5.3                                      | Northern Sweden          | Elfving 1986b                |
| 15   |                      | 70.5                                | 4.7                                      | US                       | Einspahr 1984                |
| 5  |                      | 27.0                                | 5.4                                      | Eastern Bavaria, Germany | Makeshein et al. 1989        |
| European aspen ( <i>Populus tremula</i> L.)          |                      |                                     |  |                          |                              |
| 5–24   |                      | 14–162                              | 2.9–9.1                                  | Sweden                   | Johansson 1999a              |
| 26–91  |                      | 28–501                              | 1.2–7.0                                  | Sweden                   | Johansson 2002               |
| 12   |                      | 170                                 | 14.2                                     | Finland                  | Paavilainen 1981             |
| Common alder ( <i>Alnus glutinosa</i> (L.) Gaertner) |                      |                                     |  |                          |                              |
| 4–36   |                      | 5–140                               | 0.5–7.7                                  | Sweden                   | Johansson 1999               |
| 21–91  |                      | 73–257                              | 1.5–6.1                                  | Sweden                   | Johansson 1999               |
| Grey alder ( <i>Alnus incana</i> (L.) Moench)        |                      |                                     |  |                          |                              |
| 5–35   |                      | 3–140                               | 0.7–9.9                                  | Sweden                   | Johansson 1999               |
| 21–66  |                      | 82–218                              | 2.1–5.5                                  | Sweden                   | Johansson 1999               |
| 5  |                      | 15.9                                | 6.4                                      | Estonia                  | Uri et al. 2002              |
| 19   |                      | 85                                  | 4.5                                      | Finland                  | Hytönen and Saarsalmi 2009   |
| Downy birch ( <i>Betula pubescens</i> Ehrh.)         |                      |                                     |  |                          |                              |
| 6–20   |                      | 7–61                                | 0.6–5.1                                  | Sweden                   | Johansson 1999               |
| 19   |                      | 112                                 | 5.9                                      | Finland                  | Hytönen and Saarsalmi 2009   |
| 20   |                      | 35                                  | 1.7                                      | Finland                  | Mälikönen and Saarsalmi 1982 |
| Silver birch ( <i>Betula pendula</i> Roth)           |                      |                                     |  |                          |                              |
| 8–32   |                      | 6–175                               | 0.7–8.44                                 | Sweden                   | Johansson 1999               |
| 6  |                      | 14                                  | 2.3                                      | Sweden                   | Telenius 1999                |
| 14   |                      | 59                                  | 4.2                                      | Finland                  | Ferm and Kaunisto 1983       |
| 19   |                      | 108                                 | 5.7                                      | Finland                  | Hytönen and Saarsalmi 2009   |
| Hybrid poplar  |                      |                                     |  |                          |                              |
| 6  |                      | 45                                  | 7.5                                      | Sweden                   | Telenius 1999                |
| 4–73   |                      | 19–438                              | 3–15                                     | Sweden                   | Johansson and Karačić 2011   |

The mean leaf area index (LAI) for hybrid aspen in the present study was  $2.24 \pm 0.58$  (0.68–3.35). There are few LAI values reported for hybrid aspen but Johansson (2002) reported

$2.45 \pm 0.66$  (1.80–3.19) for 26 to 91-year-old European aspens (*Populus tremula* L.). Tadaki (1966) reported that a normal range of LAI for aspen is 2.6 (2–5). Pollard (1972) reported LAI of 2.4,

2.9 and 1.9 respectively on 5-, 15- and 52-year-old stands of trembling aspen (*Populus tremuloides* Michx.) growing in the Ottawa Valley (lat. 46°00' N. and long. 77°26' W.). Peterson et al. (1970) reported a LAI of 1.8 for trembling aspen growing in Alberta. According to Korner (1991) SLA varies greatly among species and is influenced by environmental conditions. In addition, SLA is an important leaf characteristic, since it is positively related to net assimilation rate (Korner 1991). The SLA for hybrid aspen in the present study was  $161.7 \pm 81.1$  ( $33\text{--}351$ )  $\text{cm}^2 \cdot \text{g}^{-1}$  d.w. Tullus et al. (2007) reported SLA-values of  $102 \text{ cm}^2 \cdot \text{g}^{-1}$  in a study of 5-year-old hybrid aspen stands in Estonia. Yu (2001) reported  $137 \text{ cm}^2 \cdot \text{g}^{-1}$  for a 5-year-old hybrid aspen clone trial. PLA is used for light interception studies. According to Ridge et al. (1986), Dunlap and Stettler (1998) and Taylor et al. (2001) the leaf size is a good indicator of biomass productivity in fast growing trees. In the present study the mean projected leaf area (PLA) for hybrid aspen leaf area was  $48.9 \pm 13.8 \text{ cm}^2$  range 30–71. In the study by Tullus et al. (2007) PLA was  $19 \text{ cm}^2$  for 5-year-old hybrid aspens and in a study by Yu (2001) PLA was  $18 \text{ cm}^2$ .

In spite of these stands were not managed and damaged by wild habitat mean MAI was  $7$  ( $3\text{--}13$ )  $\text{t} \cdot \text{ha}^{-1} \cdot \text{a}^{-1}$ . MAI values of  $>10 \text{ t} \cdot \text{ha}^{-1} \cdot \text{year}^{-1}$  are possible for hybrid aspen plantations, and rotation periods of 10–20 years have been shown to be economic and practical (Benson and Einspahr 1967). Most hybrid aspen plantations growing on farmland are located near populated areas. Therefore, transport distances tend to be short for these harvested biomass fuels. The biomass used for bioenergy can be sold to district heating plants or to neighbors using biomass as a fuel. After harvesting the stand, a lot of suckers ( $50,000\text{--}100,000$  suckers  $\cdot \text{ha}^{-1}$ ) will emerge (Rytter 2006). The biomass production of 4-year-old 57000 suckers  $\text{ha}^{-1}$  was  $39 \text{ t} \cdot \text{ha}^{-1}$  corresponding to  $\text{MAI} = 9.5 \text{ t} \cdot \text{ha}^{-1} \cdot \text{a}^{-1}$ . Small trees, such as 5–10 year-old suckers, can be harvested using modern machines and then chipped. Alternatively, such trees can be felled using a chainsaw or whole-tree harvesters.

## Conclusions

In the hybrid aspen stands examined in the present study biomass production was  $135 \pm 53 \text{ t} \cdot \text{ha}^{-1}$  with a range of  $42\text{--}219 \text{ t} \cdot \text{ha}^{-1}$ . The average basal area was  $24.6 \pm 10.0 \text{ m}^2 \cdot \text{ha}^{-1}$  ( $7.1\text{--}42.6$ ). The stands were mostly not managed and in some stands there were low number of stems per hectare caused by damages of wild habitats and poor initial soil preparations before planting. These findings indicate the need of proper soil treatment before planting and an efficient fence, which must frequently be observed. Overall the results indicate that there is a high potential for using hybrid aspen stands in Sweden for viable biomass production.

## Acknowledgements

E. Temnerud, L. Hedman, L., B. Hjelm and M. Cardoso measured and cut stands. Miss M. Johansson weighed and dried the tree components, carried out the tree ring analyses and the leaf characteristic measurements in the laboratory. Linguistic revision

was made by Sees-Editing Ltd UK. The entire above are gratefully acknowledged. Skogssällskapet foundation provided financial support.

## References

- Andersson E, Tuimala A. 1980. Definitioner, mm, vid vedundersökningar. (Definitions for Wood Technology). The Nordic Joint Group in Wood Technology Research Note No. 4: 53–66. (in Swedish)
- Benson MK, Einspahr DW. 1967. Early growth of diploid and triploid hybrid aspen. *Forest Science*, **13**(2): 150–155.
- Bolstad PV, Gower ST. 1990. Estimation of leaf area index in fourteen southern Wisconsin forest stands using a portable radiometer. *Tree Physiology*, **7**: 115–24.
- Bréda NJJ. 2003. Ground-based measurements of leaf area index: a review of methods, instruments and current controversies. *Journal of Experimental Botany*, **54**(392): 2403–2417.
- Christersson L. 1996. Future research on hybrid aspen and hybrid poplar cultivation in Sweden. *Biomass & Bioenergy*, **11**: 109–113.
- Dickmann DJ. 2001. An overview of genus *Populus*. In: D. I. Dickmann, J.G. Icebrands and J. E. Eckenwalder, J. Richardson (eds.), *Poplar culture in North America*. Ottawa: NRC Research Press, pp. 1–42.
- Dunlap JM, Stettler RF. 1998. Genetic variation and productivity of *Populus trichocarpa* and its hybrids. X. Trait correlations in young black cottonwood from four river valleys in Washington. *Trees-Structure and Function*, **13**: 28–39.
- Einspahr DW. 1984. Production and utilization of triploid hybrid aspen. *Iowa State Journal of Research*, **58**: 401–409.
- Einspahr DW, Wyckoff GW. 1975. Aspen hybrids promise future source of Lake States fiber. *Pulp and Paper*, **49**(12): 118–119.
- Einspahr DW, Benson MK, Peckham JR. 1968. In: Proceedings of the Eighth Lake States Forest Tree Improvement Conference, Research Paper NC-23. St. Paul, Mn. U.S. Forest Service. North Central forest Experiment Station, pp. 24–27.
- Ekström G. 1926. Klassifikation av svenska åkerjordar. (Classification of Swedish soil types on farmland). SGU. Serie C. Avhandlingar och Uppsatser, 345: p. 161. (in Swedish)
- Elfving B. 1986a. Ett försök med åkerplantering av hybridasp och gran nära Sundsvall. (A trial with a plantation of hybrid aspen on abandoned farmland close to the city of Sundsvall). SST, 5: pp. 43–45. (In Swedish).
- Elfving B. 1986b. Odlingvärdet av björk, asp och al på nedlagd jordbruksmark i Sydsverige. Summary: The value of growing birch, aspen and alder on abandoned fields in southern Sweden. SST, 5: pp.31–39. (In Swedish).
- Eriksson H. 1984. Yield of aspen and poplars in Sweden. Department of Ecology and Environmental Research. Swedish University of Agricultural Sciences. Report, 15: pp. 393–419.
- Ferm A, Kaunisto S. 1983. Luontaisesti syntyneiden koivumetsiköiden maanpäällinen lehdetön biomassatuotos entisellä turpeennostoalueella kihniön aitonnesevalla. (Summary: Above-ground leafless biomass production of naturally generated birch stands in a peat cut-over area at Aitonneseva, Kihniö). *Folia Forestalia*, **558**: 1–32. (in Finnish)
- Hofmann-Schielle C, Jug A, Makeschin F, Rehfuess FE. 1999. Short-rotation plantations of balsam poplars, aspen and willows on former arable land in the Federal Republic of Germany. I. Site - growth relationships. *Forest*



- Ecology and Management*, **121**: 41–55.
- Hytönen J, Saarsalmi A. 2009. Long-term biomass production and nutrient uptake of birch, alder and willow plantations on cut-away peatland. *Biomass and Bioenergy*, **33** (9): 1197–1211.
- Ilstedt B, Gullberg U. 1993. Genetic variation in a 26-year old hybrid aspen trial in Southern Sweden. *Scandinavian Journal of Forest Research*, **8**: 185–192.
- Jacobsen B. 1976. Hybridasp (*Populus tremula* x *Populus tremuloides* Michx.). (Summary: Hybrid aspen (*Populus tremula* x *Populus tremuloides* Michx.)). *Det Forstlige Forsøgsvæsen i Danmark*, **34**: 317–338. (In Danish)
- Johansson T. 1996. Estimation of canopy density and irradiance in 20- to 40-year-old birch stands (*Betula pubescens* Ehrh. and *Betula pendula* Roth). *Trees*, **10**: 223–230.
- Johansson T. 1999a. Biomass equations for determining fractions of European aspen (*Populus tremula* L.) growing on abandoned farmland and some practical implications. *Biomass & Bioenergy*, **17**: 471–480.
- Johansson T. 1999b. Biomass equations for determining fractions of pendula and pubescent birches growing on abandoned farmland and some practical implications. *Biomass & Bioenergy*, **16**: 225–238.
- Johansson T. 1999c. Dry matter amounts and increment in 21- to 91-year-old common alder and grey alders and some practical implications. *Canadian Journal of Forest Research*, **29**: 1679–1690.
- Johansson T. 1999d. Site index curves for common and grey alders growing on different types of forest soil in Sweden. *Biomass & Bioenergy*, **18**: 147–159.
- Johansson T. 2000. Biomass equations for determining fractions of common and grey alders growing on abandoned farmland and some practical implications. *Biomass & Bioenergy*, **18**: 147–159.
- Johansson T. 2002. Increment and biomass in 26- to 91-year-old European aspen and some practical implications. *Biomass & Bioenergy*, **23**: 245–255.
- Johansson T. 2006. Site index conversion equations *Picea abies* and five broadleaved species in Sweden: *Alnus glutinosa*, *Alnus incana*, *Betula pendula*, *Betula pubescens* and *Populus tremula*. *Scandinavian Journal of Forest Research*, **21**: 14–19.
- Johansson T. 2008. Sprouting ability and biomass production of downy and silver birch stumps of different diameters. *Biomass & Bioenergy*, **32**: 944–951.
- Johansson T. 2011. Site index curves for poplar growing on former in Sweden. *Scandinavian Journal of Forest Research*, **26**: 161–170.
- Johansson T, Karačić A. 2011. Increment and biomass in hybrid poplar and some practical implications. *Biomass & Bioenergy*, **35**: 1925–1934.
- Johansson H. 1953. Hybridaspens ungdomsutveckling och ett försök till framtidsprognos. (Growth of young hybrid aspen and an attempt for future prognosis). *Svenska Skogsvårdsföreningens Tidskrift*, **51**: 73–96. (in Swedish)
- Johansson H. 1976. Das Produktionspotential der Hybridasp (*Populus tremula* x *Populus tremuloides*) in Südschweden. *Die Holzsucht*, **11/76**: 19–22. (In German)
- Karačić A, Verwijst T, Weih M. 2003. Above-ground woody biomass production of short-rotation *Populus* plantations on agricultural land in Sweden. *Scandinavian Journal of Forest Research*, **18**: 427–437.
- Karlsson B. 1991. Klonskogbruk med hybridasp. (Clone forestry by hybrid aspen). The Forestry Institute of Sweden, **3**: pp 1–4. (in Swedish)
- Kittredge J. 1944. Estimation of the amount of foliage of trees and stands. *Journal of Forestry*, **42**: 905–912.
- Korner CH. 1991. Some often overlooked plant characteristics as determinants of plant growth: a reconsideration. *Functional Ecology*, **5**: 162–173.
- Langhammer A. 1973. Et forsøk med hybridosp i Norge. (A trial with hybrid aspen in Norway). *Meldinger fra Norges Landbrukshøgskole*, **52**(6): 1–36. (in Norwegian)
- Li B, Howe GT, Wu R. 1998. Developmental factors responsible for heterosis in aspen hybrids (*Populus tremuloides* x *Populus tremula*). *Tree Physiology*, **18**: 29–36.
- Liesebach M, von Wuehlisch G, Muhs H-J. 1999. Aspen for short-rotation coppice plantations on agricultural sites in Germany: Effects of spacing and rotation time on growth and biomass production of aspen progenies. *Forest Ecology and Management*, **121**: 25–39.
- Madgwick HAI. 1971. The accuracy and precision of estimates of the dry matter in stems, branches and foliage in an old field *Pinus virginiana* stand, In: H. E. Young (ed), *Forest biomass studies*, Orono: Univ. Maine Press, US. pp.106–112.
- Madgwick HAI. 1991. Estimation stand weight – the importance of sample selection. *New Zealand Journal of Forest Science*, **21**(2/3): 180–184.
- Makeshein F, Rehfuess KE, Rüsche I, Schörry R. 1989. Anbau von Pappeln und Weiden im Kurzumtrieb auf einem ehemaligen Acker: Standortliche Voraussetzungen.
- Mäkönen E, Saarsalmi A. 1982. Hienskoivikon biomassatotos ja raviteiden mentys kokopuun korjuussa. (Summary: Biomass production and nutrient removal in whole tree harvesting of birch stands). *Folia Forestalia*, **534**: 17.
- Paavilainen E. 1981. Biomass yields and management of natural coppice stands. IEA/FE Report. NE, 19: p.50.
- Payandeh B. 1981. Choosing regression models for biomass prediction equation. *The Forestry Chronicle*, **57**(5): 229–232.
- Persson A. 1973. Ett försök med snabbväxande *Populus*-arter. (A trial with fast-growing *Populus* species). Institutionen för skogsproduktion, Skogshögskolan, Stockholm, Rapporter och Uppsatser, Nr. 27: pp.1–34. (in Swedish)
- Peterson EB, Chan YH. 1970. Aboveground standing crop, leaf area, and caloric value in an aspen clone near Calgary, Alberta. *Canadian Journal of Botany*, **48**: 1459–1469.
- Pollard DFW. 1972. Above-ground dry matter production in three stands of trembling aspen. *Canadian Journal of Forest Research*, **2**: 27–33.
- Ridge CR, Hinckley TM, Stettler RE, van Volkenburgh E. 1986. Leaf growth characteristics of fast-growing poplar hybrids *Populus trichocarpa* x *P. Deltoides*. *Tree Physiology*, **1**: 209–216.
- Rytter L. 2002. Nutrient content in stems of hybrid aspen as affected by tree age and tree size, and nutrient removal with harvest. *Biomass & Bioenergy*, **23**: 13–25.
- Rytter L. 2006. A management regime for hybrid aspen stands combining conventional forestry techniques with early biomass harvests to exploit their rapid early growth. *Forest Ecology and Management*, **236**: 422–426.
- Rytter L, Stener L-G. 2003. Clonal variation in nutrient content in woody biomass of hybrid aspen (*Populus tremula* L. x *Populus tremuloides* Michx.). *Silva Fennica*, **37**(3): 313–324.
- Rytter L, Stener L-G. 2005. Productivity and thinning effects in hybrid aspen (*Populus tremula* x *P. tremuloides* Michx.) stands in southern Sweden. *Forestry*, **78**(3): 285–295.
- SAS Institute Inc. 2006. *SAS/STAT User's Guide* Version 9.1. Cary, NC: SAS Institute Inc
- Satoo T, Madgwick HAI. 1985. *Forest biomass*. London: Martinus Nijhoff/DR W. Junk Publishers, p. 23.
- Stener L-G. 1998. Analys av fiberegenskaper för kloner av hybridasp. (Analy-

- sis of fiber quality for hybrid aspen clones), SkogForsk. Uppsala. Arbetsrapport, 387; p. 11. (in Swedish)
- Stener L-G, Karlsson B. 2004. Improvement of *Populus tremula* × *P. tremuloides* by phenotypic selection and clonal testing. *Forest Genetics*, **11**: 13–27.
- Tadaki Y. 1966. Some discussion on the leaf biomass of forest stands and trees, Bulletin Government Forest Experimental Station, Tokyo. 184: 135–162.
- Taylor G, Beckett KP, Robinson KM, Stiles K, Rae AM. 2001. Identifying QTL for yield in UK biomass poplar, Asp. *Applied Biology*, **65**: 173–182.
- Telenius BF. 1999. Stand growth of deciduous pioneer tree species on fertile agricultural land in southern Sweden. *Biomass & Bioenergy*, **16**: 13–23.
- Tullus A, Rytter L, Tullus T, Weih M, Tullus H. 2012. Short-rotation forestry with hybrid aspen (*Populus tremula* L. × *Populus tremuloides* Michx.). *Scandinavian Journal of Forest Research*, **27**(1): 10–29.
- Tullus A, Tullus H, Vares A. 2007. Early growth of hybrid aspen (*Populus* × *Wettsteinii* Hämet-Ahti) plantation on former agricultural lands in Estonia. *Forest Ecology and Management*, **245**: 118–129.
- Tullus A, Tullus H, Soo T, Pärn L. 2009. Above-ground biomass characteristics of young hybrid aspen (*Populus tremula* L. × *Populus tremuloides* Michx.) plantations on former agricultural land in Estonia. *Biomass and Bioenergy*, **33**: 1617–1625.
- Uri V, Tullus H, Lõhmus K. 2002. Biomass production and nutrient accumulation in short-rotation grey alder (*Alnus incana* (L.) Moench) plantation on abandoned agricultural land. *Forest Ecology and Management*, **161**(1-3): 169–179.
- Wettstein W. 1933. Die Kreuzungsmethode und die Beschreibung von F1 Bastarden by Populus. *Z. Züchtung. A. Pflanzenzüchtung*, **18**: 97–626. (in German)
- Woods KD, Peiveson AH, Botkin DB. 1991. Statistical error analysis for biomass density and leaf area index estimation. *Canadian Journal of Forest Research*, **21**: 974–989.
- Yu Q. 2001. Can physiological and anatomical characters be used for selecting high yielding hybrid aspen clones? *Silva Fennica*, **35**(2): 137–146.
- Yu Q, Tigerstedt PMA, Haapanen M. 2001. Growth and phenology of hybrid aspen clones (*Populus tremula* L. × *Populus tremuloides* Michx.) *Silva Fennica*, **35**(1): 15–25.
- Zar JH. 1999. *Biostatistical analysis*. Englewood Cliffs, N.J: Prentice-Hall. Inc.